

Spatial Prediction Using Combined Sources of Data

Nancy J. McMillan,¹ David M. Holland,² Michele Morara,¹ and Gregory S. Young¹

(1) Battelle, Measurement and Data Analysis Sciences, Columbus, OH
(2) U.S. Environmental Protection Agency, Office of Research and Development, National Exposure Research Laboratory, Research Triangle Park, NC

Objectives and Methods

Objectives

- Provide daily particulate matter (PM_{2.5}) and ozone (O₃) spatial surfaces for Environmental Health Tracking
- Combined predictions can be used for modeling air quality – public health relationships in the Public Health Air Surveillance Evaluation (PHASE) project
- Determine air quality non-attainment areas

Data Sources

- 24-hr average PM_{2.5} data from EPA's FRM fine Particulate Network
- Daily 8-hr maximum O₃ concentrations from the NAMS/SLAMS Network
- Community Multi-Scale Air Quality (CMAQ) daily PM_{2.5} and 8-hr maximum O₃ output over 36 km grid
- MODIS Satellite Aerosol Optical Depth (AOD) data over 10 km grid
- Eta Data Assimilation System (EDAS) meteorological data over 80 km grid
- LandScan daytime population density data
- 24-hr average PM_{2.5} data from EPA's Speciation Trends Network (STN) and Interagency Monitoring of Protected Visual Environments (IMPROVE) data for validation
- Daily 8-hr maximum O₃ concentrations from EPA's Clean Air Status and Trends Network (CASTNet) for validation





Figure 1. Predicted summer average PM_{2.5} (µg/m³) surface – Combined model (left) versus interpolated monitoring data (right)




Figure 2. Predicted O₃ (ppb) seasonal average surface – Combined model (left) versus interpolated monitoring data (right)

Methods

- Monitoring data, CMAQ output, and MODIS data can be used simultaneously to predict daily pollutant surfaces
 - Air quality monitoring data is spatially sparse, temporarily rich
 - Numerical model output has high spatial and temporal resolution, but potential for location dependent bias
 - Satellite data has high spatial and temporal resolution, but potential for location specific bias and significant missing data (cloud cover)
- Leads to more accurate predictions and prediction errors
- Draw on strengths of each data source:
 - Give more weight to accurate monitoring data in areas where monitoring data exists
 - Rely on bias adjusted model output and satellite data in non-monitored areas
- Model underlying spatial dependence and measurement errors of each data source – no blind combining
 - Monitor data
 - $X_i^j(s_{ij}) | W_i(s_{ij}), \sigma_x^2 \sim N(W_i(s_{ij}), \sigma_x^2)$
 - CMAQ Data
 - $Y_i^j(s_{ij}) | W_i(s_{ij}), \beta_{0i}, \sigma_y^2 \sim N(W_i(s_{ij}) + D_i(s_{ij})\beta_{0i}, \sigma_y^2)$
 - AOD Data
 - $S_i^j(s_{ij}) | \eta_i, V_i(s_{ij}), \sigma_s^2 \sim N(\eta_i + V_i(s_{ij}), \sigma_s^2)$
 - Underlying air quality process
 - $W_i(s_{ij}) = \mu + A_i(s_{ij})\beta_A + Z_i(s_{ij})$
 - Air quality process residuals and underlying AOD process includes an auto-regressive temporal component and a conditionally auto-regressive spatial component
 - $V_i | \sigma_v^2, \rho_v \sim N(0, \sigma_v^2(A_v^{-1}(\rho_v) \otimes A_v^{-1})^{-1})$
 - $Z_i | \sigma_z^2, \rho_z \sim N(0, \sigma_z^2(A_z^{-1}(\rho_z) \otimes A_z^{-1})^{-1})$
- Hierarchical Bayesian statistical modeling based on custom-designed Monte Carlo Markov Chain software

Table 1. Root Mean Squared Prediction Errors (RMSPE) Using Three Prediction Surfaces

Prediction Surface	O ₃			PM _{2.5}		
	Overall RMSPE	Combined Model Improvement (%)		Overall Estimate	Combined Model Improvement (%)	
		Sites 11	Days 245		Sites 60	Days 78
CMAQ Output	0.525	91%	64%	0.277	90%	87%
Kriged Monitor Data	0.521	91%	57%	0.165	58%	69%
Bayesian Combined	0.501			0.127		

Results and Future Work

Results

- Combined approach provides reliable information about the true PM_{2.5} and O₃ surfaces.
- How well does the combined approach predict to NON-MONITORED locations? Answer: Validate the model against data not used in fitting the model, use IMPROVE and STN PM_{2.5} and CASTNet O₃ data.
 - Calculate root mean squared prediction error (RMSPE)
 - Compare predictive results of combined approach to ordinary kriging

Future Work

- Use 12 km CMAQ gridded output and compare to current results with 36 km output
- Use model to refine definition of pollution non-attainment areas




Figure 3. Predictive performance of combined model versus interpolated monitoring data for PM_{2.5} (left) and O₃ (right)

Notice: Although this work was reviewed by EPA and approved for publication, it may not necessarily reflect official Agency policy. Mention of trade names or commercial products does not constitute endorsement or recommendation by EPA for use.

126lcb05db.SF-Holland



epascienceforum
Collaborative Science
for Environmental Solutions



epa.gov/scienceforum
2005